

carotid plaque surface (smooth, irregular, ulcerated), NIHSS group score before surgery (0 or TIA, 1-3, 4-6,  $\geq 7$ ) and presence of ischemic lesions at preoperative CT scans for analysis. Significance threshold was set at  $P < .05$ .

**Results:** Eighty-three patients out of 203 presented preoperative ischemic brain lesions (40.9%). In 48.3% of patients an irregular plaque surface was recorded, in 41.8% a ulcerated surface and in 9.9% a smooth surface. At between NIHSS groups analysis ulcerated plaque surface was more frequent in TIA patients ( $P = .05$ ). In those patients the presence of ulcer in the plaque surface was strongly related to absence of ischemic brain lesions ( $P = .05$ ). In stroke patients (NIHSS  $> 0$ ) the ulcerated or irregular plaque surface was slightly significantly associated to presence of preoperative brain ischemic lesions, particularly in NIHSS 1-3 group patients ( $P = .08$ ).

**Conclusions:** Aside from stenosis percentage, neurological transient and permanent brain symptoms are highly related to carotid plaque surface. Ulcerated carotid plaque surface can be responsible for plaque's micro-debris embolization or fresh thrombus formation whose brain damage mechanisms can lead to different neurological symptoms and CT scans findings.

**Author Disclosures:** L. Capoccia: Nothing to disclose; A. Esposito: Nothing to disclose; P. Fiorani: Nothing to disclose; I. P. Renzi: Nothing to disclose; E. Sbarigia: Nothing to disclose; F. Speziale: Nothing to disclose.

#### PS54.

##### Risk Scoring System to Predict Three-Year Survival in Patients Treated for Asymptomatic Carotid Stenosis

Francisco Alcocer, Marjan Mujib, Jeffrey Horn, Marc A. Passman, Mark A. Patterson, Thomas C. Matthews, William D. Jordan. Surgery, University of Alabama at Birmingham, Birmingham, AL

**Objectives:** To identify risk factors compromising 3 years survival in treated patients with asymptomatic carotid disease in complying with recently updated guidelines from the Society for Vascular Surgery.

**Methods:** Outcomes of 504 patients who underwent carotid intervention for asymptomatic carotid disease in 10 years (1999-2008) were analyzed. Hospital computerized medical records were reviewed. Social Security Death Index was queried for mortality. Patients lost to follow up before 3 years post procedure were excluded. After multivariable Cox regression analysis was done, a score was assigned for each risk factor (RF): hazard ratio (HR) 1.5-2.0 = RF 1; HR 2.1-3.0 = RF 2; HR 3.1-4.0 = RF 3. Then a Kaplan Meier plot analyzed survival differences.

**Results:** Fifteen percent of the patients did not survive beyond 3 years post procedure. Age  $> 80$  years (HR 2.0,  $P = .01$ , RF = 1), Chronic obstructive pulmonary disease (HR 3.6,  $P = .001$ , RF = 3), Chronic kidney disease (CKD) stage 3 (HR 2.0  $P = .07$ , RF = 1), CKD stage 4 (HR 3.3,  $P = .04$ , RF = 3), Coronary artery disease (HR 2.46,  $P = .001$ , RF = 2) and Diabetes Mellitus (HR 1.8,  $P = .02$ , RF = 1) negatively influenced 3 year survival.




Fig. Kaplan-Meier Survival Plots by Risk Score

Thirty days mortality rate was 0.9% and was not influenced by score values; however, a cumulative score  $< 3$  was associated with 3 year survival of 93%.

**Conclusions:** Despite low 30 day mortality rate, increasing risk factors are associated with decreased 3 year survival. This finding may impact clinical decision for management of asymptomatic carotid disease.

**Author Disclosures:** F. Alcocer: Nothing to disclose; J. Horn: Nothing to disclose; W. D. Jordan: Nothing to disclose; T. C. Matthews: Nothing to disclose; M. Mujib: Nothing to disclose; M. A. Passman: Nothing to disclose; M. A. Patterson: Nothing to disclose.

#### C9d: Poster Session - Cerebrovascular including Great Vessels (2)

#### PS56.

##### Impact of Distal Protection Filter Design in Thirty-Day Outcomes of Carotid Artery Stenting

Natasha Loghmanpour<sup>1</sup>, Gail Siewiorek<sup>1</sup>, Kelly Wanamaker<sup>2</sup>, Mark Wholey<sup>3</sup>, Rabih Chaer<sup>3</sup>, Satish Muluk<sup>2</sup>, Ender A. Finol<sup>4</sup>. <sup>1</sup>Carnegie Mellon University, Pittsburgh, PA; <sup>2</sup>West Penn Allegheny Health System, Pittsburgh, PA; <sup>3</sup>The University of Pittsburgh Medical Center, Pittsburgh, PA; <sup>4</sup>The University of Texas at San Antonio, San Antonio, TX

**Objectives:** To review retrospectively records of patients treated with carotid artery stenting (CAS) to investigate correlations between clinical variables, distal protection filter (DPF) characteristics, and 30-day peri/post-procedural outcomes.

**Methods:** This is a multicenter, single-arm study of DPF-protected CAS in the Pittsburgh region between 2000 and 2011. Analysis of peri/post-procedure complications included myocardial infarction (MI), transient ischemic attacks (TIA), stroke, death, and a composite of all adverse events (AE). Characteristics for the DPFs were previously determined in vitro and were used to find correlations with CAS outcomes. Univariate, multivariate, and goodness-of-fit analyses were performed.

**Results:** 731 CAS procedures employing six different DPF were analyzed. Peri/post-procedural AE included 19 TIAs (2.6%), 38 strokes (5.2%), 1 MI (.1%), 19 deaths